

Special Local Need Registration - 24(c)

SUPPLEMENTAL LABELING FOR DISTRIBUTION AND USE ONLY IN OHIO

Stratego™

Fungicide

EPA Reg. No. 3125-562

EPA SLN No. OH-02001

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions and precautions on the EPA-registered label for all products used are to be followed.

This labeling must be in the possession of the user at time of pesticide application.

Wheat

Stratego will provide effective control of wheat rusts (*Puccinia graminus*, *P. recondita*, *P. striiformis*), powdery mildew (*Erysiphe graminis*), and Stagonospora leaf and glume blotch. (*Stagonospora nodorum*).

Applications may be either by ground equipment or aerially. Stratego may be applied at any growth stage through Feekes Growth Stage 10.5 (full head emergence). Do not apply after this growth stage to avoid possible illegal residues.

Disease	Rate	Comments
Rusts Powdery Mildew Stagonospora Leaf and Glume Blotch	10 fl. oz./A	Begin applications preventively when conditions are favorable for disease development.
Notes: To avoid possible illegal residues: 1. Do not exceed more than 10 fluid ounces of Stratego per acre per season. 2. Do not apply within 40 days of harvest. 3. Do not graze or feed livestock treated wheat, straw, forage or hay. 4. Do not cut the green crop for hay or silage. 5. After harvest, the straw may be used for bedding.		

OHSTSL01_034 03/11/02

Bayer Corporation
Crop Protection
Box 4913
Kansas City, MO 64120-0013
(800) 842-8020
<http://uscrop.bayer.com>

IMPORTANT

Before using this product, read and carefully observe the directions, cautionary statements and other information appearing on the product packaging label. This product is sold subject to the Conditions of Sale set forth on the container label.

