

**FOR DISTRIBUTION AND USE ONLY WITHIN OHIO.  
 FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED  
 LABEL. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE  
 APPLICATION.**

Read "Limit of Warranty and Liability" before using. If terms are not acceptable, return at once unopened.

The liability of UCB Chemicals Corporation ("UCB") for any injury, loss or damage to property associated with the use of this product under any theory of warranty, negligence, or strict liability shall be limited to a refund of the purchase price of the product.



**Ferbam Granuflo®**  
**Fungicide**  
**SPECIAL LOCAL NEED LABEL**

EPA Reg No. 45728-7

EPA SLN No. OH-990004

Active Ingredient:

Ferbam (Ferric dimethyldithiocarbamate).....76.0%

Inert Ingredients.....24.0%

Total.....100.0%

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique detalle.  
 (If you do not understand the label, find someone to explain it to you in detail.)**

**EMERGENCY CALLS: (800) 424-9300 OR (404) 801-3248 OR (404) 433-6421**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

CROP	DISEASE	USE RATE	USE DIRECTIONS
Caneberries (including raspberries, loganberries, youngberries, blackberries)	Rust, Spur Blight, Cane and Flower Botrytis, Leaf Spot, Anthracnose	1.5 - 3 lbs/A	<p>Apply at delayed dormant. Repeat when the blossoms are in the bud stage (when new canes are about 8 to 10 inches high). Cover new shoots thoroughly. Apply a third spray two weeks later. For post-harvest applications, apply during late summer or fall when temperatures start to decline through October 31. Additional applications can be made at 2- week intervals if weather is favorable for disease. Use the higher rate under high disease pressure.</p> <p>Application of up to 100 gallons of water per acre is recommended to obtain thorough coverage. Adjust total volume applied per acre to achieve thorough coverage of canes and foliage.</p> <p>Use of a suitable adjuvant may help improve coverage. Do not use a penetrating-type adjuvant.</p> <p>Do not apply within 40 days of harvest. Do not apply more than 15 lbs (11.4 lbs active ingredient) per crop season.</p>

**DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.**

**Limit of Warranty and Liability**

Neither UCB Chemicals Corporation or the seller makes any warranty of merchantability, fitness of purpose, or otherwise, expressed or implied, concerning use of this product in accordance with these provisions. The user and/or grower acknowledges the preceding disclaimer and accepts liability for any possible damage resulting from this use, whether under theories of warranty, negligence, or strict liability which exceed the purchase price of the product.

UCB and the seller make this product available to the user because the benefit and utility, in the opinion of the grower and/or user, outweigh the extent of potential crop injury, loss, or damage associated with use of the product.



Chemicals Corporation  
 2000 Lake Park Drive  
 Smyrna, Georgia 30080  
 800/426-3820

Granuflo® is a Registered Trademark of UCB S.A., a Belgian corporation.