

syngenta

FIFRA

Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN
THE STATE OF OHIO**

RIDOMIL GOLD[®] SL

EPA Reg. No. 100-1202

EPA SLN No. OH-110005

**Use of Ridomil Gold SL in Tobacco Transplant Water for Control of
Black Shank**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR PEST CONTROL, ILLEGAL RESIDUES, OR CROP INJURY.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Ridomil Gold SL label.

This label must be in the possession of the user at the time of application.

Specific Use Directions in Tobacco

Use in Transplant Water:

- Apply in transplant furrow while planting tobacco seedlings for control of black shank caused by *Phytophthora nicotianae*
- Apply 4-8 oz/A (0.25-0.5 pt/A) in at least 200 gallons of transplant water per acre.
- Use the high rate if the disease epidemic is expected to be severe.
- Make at least 1 subsequent application of Ridomil Gold SL at first cultivation and/or layby if necessary, for control of black shank.

Precautions

- There is a risk of plant injury with transplant water application of Ridomil Gold SL, especially when applied in less than 200 gallons carrier volume/acre. Crop injury is most likely when application is in less than 100 gallons /acre. Injury is temporary and should disappear within 3 weeks.
- Do not apply in less than 200 gallons for burley and dark tobacco.
- Pre-mixing Ridomil Gold SL in a tank separate from the transplant water source tank will help to prevent negative interactions with fertilizers or other pesticides in the transplant water solutions.
- Do not apply to stressed seedlings, or during hot and dry conditions due to injury potential.
- A variety with resistance to race 0 and race 1 of the black shank pathogen should be planted

Restrictions

- Do not exceed the equivalent of 1.5 lb. a.i./A (3 pts/A) per crop of mefenoxam containing products.

Ridomil Gold® trademark of a Syngenta Group Company

24(c) registrant:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

Label Code: OH1202017AA0411